

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

JC05 Rec'd PCT/PTO 25 MAR 2002

Attorney's Docket Number
18377-0012

U.S. Application No.
(if known, see 37 CFR 1.5)

10/089130

International Application No.
PCT/AU00/01184

International Filing Date
22 September 2000

Priority Date Claimed
24 September 1999

Title of Invention

RECOGNITION OF DIFFERENCES IN CELL CYCLE STRUCTURE BETWEEN STEM
AND DIFFERENTIATED CELLS

Applicant(s) for DO/EO/US

RATHJEN, Peter David and DALTON, Stephen

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), and (9) and (21) indicated below.
4. ☐ The US has been elected by the expiration of 19th month from the earliest claimed priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is attached herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ An English language translation of the International Application into English (35 U.S.C. 371(c)(2))
 - a. ☐ is attached herewith.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
 - c. ☒ translation not required as the application was filed in English.
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 20. below concern document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.
14. ☐ A SECOND or SUBSEQUENT preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
18. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☐ Other items or information:

Express Mail Label No. EL690570248US

Date: March 25, 2002

Page 1 of 2

ATTACH CUSTOMER BAR CODE LABEL BELOW



29052

PATENT TRADEMARK OFFICE

U.S. Application No. 10/089130 <small>(37 CFR 1.53)</small>	International Application No. PCT/AU00/01184	Attorney's Docket Number 18377-0012
---	---	--

21. ☒ The following fees are submitted: CALCULATIONS PTO USE ONLY

BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):

Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO.. **\$1040.00**

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO **\$890.00**

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO **\$740.00**

International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) **\$710.00**

International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) **\$100.00**

ENTER APPROPRIATE BASIC FEE AMOUNT =				\$ 1,040.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
Claims	Number Filed	Number Extra	Rate		
Total claims	108 - 20 =	88	x 18.00	\$ 1,584.00	
Independent Claims	8 - 3 =	5	x 84.00	\$ 420.00	
Multiple Dependent Claims (if applicable)				+ 280.00	\$ 280.00
TOTAL OF ABOVE CALCULATIONS =				\$ 3,324.00	
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				-	\$ 1,662.00
SUBTOTAL =				\$ 1,662.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$ 1,662.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				+	
TOTAL FEES ENCLOSED =				\$ 1,662.00	
				Amount to be refunded:	\$
				charged:	\$

a. ☒ A check in the amount of **\$1,662.00** to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. 19-5029 in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment, to Deposit Account No. 19-5029. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:


William L. Warren, Esq.

SUTHERLAND ASBILL & BRENNAN, LLP

999 Peachtree Street, N.E.

Atlanta, Georgia 30309

Telephone: 404-853-8000


 William L. Warren, Reg. No. 36,714

FORM PTO-1390 (Rev. 11-2000) adapted Page 2 of 2

Recognition of differences in cell cycle structure between stem and differentiated cells

The present invention relates to methods for using molecules relating to the control of the cell cycle, and cell proliferation, to improve technology relating to pluripotent, multipotent and differentiated cells. More particularly, the present invention relates to methods for identifying pluripotent cells and partially differentiated cells, to methods for enhancing the maintenance and proliferation of pluripotent cells and partially differentiated cells, to methods for isolating new populations of pluripotent cells and partially differentiated cells, and facilitating their maintenance and proliferation *in vitro*, to methods for reprogramming of differentiated somatic cells so that the cells are converted to a less differentiated state, including to a state of pluripotency or multipotency, and to methods for selecting dedifferentiated cells, including those derived by reversion of differentiated or partially differentiated cells, in a mixed population of cells comprised of differentiated and dedifferentiated cells.

This invention also relates to methods for regulating the differentiation of cells, including pluripotent cells and multipotent cells, and to methods for prolonging the lifespan *in vitro* of pluripotent, multipotent or differentiated cells. Also within the scope of the present invention are cells, embryos and animals produced using the methods referred to above. In addition, uses of the cells, embryos and animals derived by these methods are within the scope of this invention.

In this patent application the term "pluripotent" refers to cells that can contribute substantially to all tissues of the developing embryo. "Multipotent" or "partially differentiated" refers to partially differentiated cells that are able to differentiate further into more than one terminally differentiated cell type. Such cells include, but are not limited to haematopoietic stem cells and neural stem cells.

"Maintenance of pluripotent cells" is to be understood as the

maintenance of such cells *in vitro* in an undifferentiated state. It may also include, but does not always include, the understanding that these cells are immortal.

Control of the cell division cycle and roles for cell cycle regulators
5 ***in chromatin remodelling***

The cell division cycle is normally composed of four distinct phases, which in typical somatic cells take 18-24 hours to complete. The S-phase represents the period when chromosomal DNA is duplicated, this is then followed by a gap phase (G2) where cells prepare to segregate chromosomes
10 between daughter cells during M-phase. After completion of M-phase, cells enter a second gap phase, G1, which separates M- from S-phase. G1 is of special significance because it is here that a cell decides to continue dividing or withdraw from the cell cycle.

At the molecular level, the cell cycle is controlled by waves of cyclin-
15 dependent protein kinase (Cdk) activities that are activated only at specific times and which drive the cell cycle transitions by phosphorylation of specific substrates. For activity, each Cdk catalytic subunit requires a cyclin regulatory subunit. Cdks acting at the G1 phase include Cdk2 which is regulated by cyclin E, and Cdk4 and Cdk6 which are regulated by cyclin D activities. Additional
20 levels of control are provided by cyclin-dependent kinase inhibitors, such as p16.

Mitogenic signals, under normal circumstances, dictate whether a cell divides or arrests (Figure 1). The major step in mitogen-dependent control of cell proliferation comes at a point in G1, known as the Restriction Point (R-
25 point), where cells commit to entering S-phase and to another round of cell division. At the molecular level, mitogen-dependent intracellular signalling pathways control this 'proliferative switch' by activation of cyclin-associated Cdk activities, which function primarily by phosphorylating the retinoblastoma tumour suppressor protein (pRb) and its family members, including p107 and



p130. This then allows for the dissociation of pRb from E2F transcription factors, resulting in derepression of target genes essential for the G1 to S transition. On the other hand, failure to activate cyclin-associated kinase activities will leave hypo-phosphorylated pRb associated with E2F, and exit from the cell cycle will follow (Figure 1). Significantly, loss of R-point control is an important step in tumorigenesis (Hall & Peters, 1996; Palermo & Peters, 1996). Another G1 Cdk activity, consisting of Cdk2 and cyclin E, collaborates with Cdk4,6-cyclin D to fully phosphorylate pRb and to promote the G1-S transition. Under some circumstances, elevated levels of cyclin E can relieve the requirement for cyclin D-associated Cdk activities, thus bypassing the R-point (Jiang et al, 1998) and cells lacking the pRb-pathway have a reduced requirement for Cdk4-cyclin D activities (Sherr & Roberts, 1995) and can pass from G1 through to S-phase using Cdk2 activity. Another Cdk2 activity, this time associated with cyclin A, is important for progression through S-phase in typical somatic cells (Pagano et al, 1992).

Cell cycle regulation clearly involves transcriptional control through pRb and its effects on E2F transcription factors. There is also now an emerging view that the molecules effecting cell cycle regulation are also linked to chromatin remodelling and differentiation. For example the transcriptional activities of pRb are associated with changes in chromatin structure (Brehm & Kouzarides, 1999). Furthermore histone acetyl transferases have a well established role in global control of gene expression and chromatin remodelling, with consequent effects on cell cycle regulation and differentiation (Kouzarides 1999). Changes in the activities of chromatin remodelling complexes have, in fact, been heavily implicated in the processes of cellular differentiation in a wide range of mammalian cell types (Kawasaki et al, 1998). In summary cell cycle regulation, and in particular cyclin-Cdk regulation, is associated with transcriptional control and changes in chromatin structure, and is likely to play an important role in global profiles of gene expression and differentiation.

Proliferation and differentiation during development

During mammalian development the differentiation of pluripotent cells and partially differentiated cells is coordinated closely with changes in proliferation.

- 5 In the mouse, development of the embryo in the first few days following implantation is a very dynamic period characterised by rapid cell proliferation and differentiation. At the time of implantation at 5.0 days post coitum (dpc) the embryo comprises a central ball of Inner Cell Mass (ICM) cells which are pluripotent (they can give rise to all cell types of the later embryo and adult).
- 10 These are surrounded by the extraembryonic primitive endoderm and trophoctoderm lineages. Around 5.0dpc the pluripotent cells, referred to from this time as epiblast, commence a period of rapid proliferation which accompanies transition from the ICM ball of cells, to a unicellular layer of pluripotent primitive ectoderm (Figure 2). The entire embryo arises from
- 15 differentiation of the primitive ectoderm into the three germ layers ectoderm, endoderm and mesoderm during gastrulation which initiates at around 6.5dpc with the appearance of a structure called the primitive streak which forms at the posterior region of the embryo (Figure 2). Pluripotent cells migrate through the streak losing pluripotence and emerge as differentiated germ layer cells.
- 20 From this time pluripotent cells in the embryo are restricted to the future germ cells.

- Immediately following implantation the region of the embryo that will contribute to the adult, the primitive ectoderm, consists of around 30 cells, which divide once every 10-12 hours. This rate of cell division is marginally
- 25 faster than that in the intestinal crypt, the most rapidly cycling cells in the adult. The embryo maintains this rapid rate of cell division in the primitive ectoderm cells for the next 48 hours. Just before gastrulation, the cell cycle of the primitive ectoderm decreases even further to around 6 hours (Figure 2). Indeed some propose the existence of a sub-population of cells within the
- 30 primitive ectoderm with even faster cell cycle times of 3 to 4 hours. This is a

quite remarkable shortening of the cell cycle, especially when compared with the length of most somatic cell cycles, which range from 18-24 hours (Figure 2). Accelerated proliferation within the embryonic epiblast therefore precedes, or coincides with, the differentiation of primitive ectoderm and hence, loss of
5 unlimited differentiation potential associated with pluripotency.

The very rapid expansion of the primitive ectoderm generates sufficient cell numbers to support formation of the three germ layers over a short developmental time-frame. This burst of cell proliferation is considered to be the force which drives gastrulation and can account for the transformation of a
10 single layered embryo into the three germ layers, the definitive ectoderm, mesoderm and endoderm. Cell division in the newly formed mesoderm is not considered to play a significant role in expansion of this layer, nor the endoderm. Cell cycle length in the newly formed mesodermal cells has been measured at 12 hours and given that a complete mesodermal layer forms
15 within 24 hours of the onset of gastrulation, most of the increase in this germ layer must be contributed from elsewhere, namely the primitive ectoderm (Hogan et al, 1994).

In the post-gastrulation embryo it appears that all cells have cycles of >12 hours. It is therefore apparent that in the early postimplantation embryo
20 the rate of proliferation firstly increases dramatically just prior to and during gastrulation, but then slows after the cells lose pluripotency in the post-gastrulation embryo, and at some time they differentiate.

In the mammal the close association between differentiation and alterations in proliferation continues beyond embryonic development, and
25 extends throughout adult life.

Multipotent stem cell populations play a critical role in mammalian development and in normal renewal of differentiated cells. Many differentiated cells are not replaced by proliferation of existing differentiated cells, but by the

differentiation of partially differentiated (precursor) cells called multipotent stem cells. Multipotent stem cells are not terminally differentiated themselves, but are competent to differentiate into one or more terminally differentiated cell types (multipotent differentiation potential). For example the stem cell for haemopoiesis can differentiate into at least 9 different kinds of blood cells. The rates of renewal and differentiation of stem cells are coupled to control the rate of differentiated cell production and prevent depletion of the stem cell population. Multipotent stem cells have a capacity for renewal which, if not infinite, extends beyond the lifetime of the animal.

10 The changes in the rate of cellular proliferation during embryogenesis and the continued close association between differentiation and changes in proliferative rates during later development have been described previously (Hogan et al, 1994), but the cell cycle structure and molecular mechanisms underlying these changes have not been rigorously evaluated.

15 The availability of murine pluripotent cells in vitro has led to the development of powerful model systems for investigating mechanisms of early development. In particular pluripotent cells provide an opportunity to investigate the molecular events responsible for the close association between pluripotency and cell cycle characteristics, and the link between differentiation and changes in regulation of the cell cycle.

20 Murine pluripotent cells can be isolated from the preimplantation embryo and maintained in vitro as ES cells. ES cells retain pluripotence indefinitely and display the properties of stem cells, including competency to differentiate into all cell types, and the ability for indefinite self-renewal. Early primitive ectoderm-like (EPL) cells are also pluripotent stem cells. They differ in some properties to ES cells, and have the capacity to revert to ES cells in vitro. They can be derived from ES cells or other types of pluripotent cells, and are the in vitro equivalent of primitive ectoderm cells of postimplantation embryos. As such, EPL cells can also be established in vitro from cells isolated from the

primitive ectoderm of postimplantation embryos. The properties of EPL cells, factors required for their maintenance and proliferation in vitro, and their ability to differentiate uniformly in vitro to form essentially homogeneous populations of partially differentiated and differentiated cell types are described fully in PCT/AU99/00265, to applicants, the entire disclosure of which is incorporated herein by reference. Cells of the primordial gonad, primordial germ cells (PGCs), also retain pluripotency during embryonic development, and can be isolated and cultured in vitro as embryonic gonadal (EG) cells. Embryonic carcinoma (EC) cells may also be pluripotent.

While pluripotent cells and partially differentiated cells have long been recognised as ideally suited to a range of applications, in practice technical barriers have generally restricted their use in the prior art. Specific opportunities and shortcomings of stem cell technology include:

Genetic Modification of Livestock

The potential for genetic modification of livestock species for agricultural, medical and commercial application is enormous. Proof of concept has been achieved by application of traditional pronuclear injection techniques for the expression of therapeutic proteins in ruminant milk (Janne et al, 1998). However, the genetic modification of animals for applications such as xenotransplantation (Platt & Lin, 1998), has been severely limited by inherent shortcomings in pronuclear injection technology. Firstly, integration of the injected DNA is inefficient and random, leading to difficulties in achieving appropriate transgene expression. Secondly, modification of endogenous genes is possible only in exceptional circumstances.

ES cell and nuclear transfer technologies have the potential to overcome these deficiencies:

Relatively robust technologies have been developed for the creation of mice that are genetically modified in a predetermined manner (Bradley et al,

1992). Methodologies based on homologous recombination are used for efficient and precise genetic alteration of endogenous genes in ES cells, including alteration of individual nucleotides. Genetically modified ES cells are then used as a vector to transmit the genetic modification through the germline of chimaeric animals to all cells of F1 offspring.

The ability to genetically alter commercially valuable species in a precise way has been limited by the inability to generate stable, proliferating pluripotent cell lines from non-rodent species. A novel approach for improved isolation and maintenance of pluripotent cells in vitro from a wide range of species, that were previously refractory to isolation by conventional approaches would be a significant advance in the art. Widespread in vitro availability of robust pluripotent cells from commercially useful species would thereby provide an opportunity for their precise genetic manipulation and use in agricultural, commercial and medical applications.

The creation of Dolly the cloned sheep (Wilmut et al, 1997) by nuclear transfer identified a second potential route to precise genetic modification of mammals. It is envisaged that genetic modification will be carried out in primary somatic cell cultures by homologous recombination technologies developed for use in mouse ES cells. The nuclei from genetically modified cells would be used as karyoplast for nuclear transfer to create genetically altered animals. A problem with this technology is that the efficiency of nuclear transfer is low; only a small proportion of nuclear transfer embryos develop to live animals. This inefficiency is due largely to problems with reprogramming of the somatic nucleus to a pluripotent state after transfer to the recipient oocyte.


Novel Human Therapies

A second area in which stem cell technology is expected to have commercial impact is in the development of human therapeutics for gene therapy and cell-based therapy (Smith, 1998; Rathjen et al, 1998). These technologies will be used for treatment of diseases in which cell replacement is

likely to be of value (ie Parkinson's, chronic viral infection), for correction of genetic defects (gene therapy), and for delivery of protein pharmaceuticals. Alternative strategies are based on the use of somatic multipotent stem cells and embryonic stem cells. The attraction of somatic stem cells for gene
5 therapy is that, being immortal, administration of genetically altered stem cells should provide long term cures for genetic disease.

Significant problems have been encountered in the realisation of this technology. In many cases, identification and culture of stem cells for target tissues has not been achieved. Furthermore, even when cell populations can
10 be enriched, for example in the case of haemopoietic stem cells, these have proven refractory to genetic manipulation. In particular the failure of cultured stem cells to proliferate clonally in culture prevents the use of homologous recombination-based techniques for modification of endogenous genes.

Human application of ES cell technologies provides a route of great
15 promise for the development of novel human therapies (Smith, 1998; Rathjen et al, 1998). Briefly, human ES cells would be differentiated in vitro to an appropriate cell type for transplantation. The ES cells could also be genetically modified using the homologous recombination based technologies prior to controlled differentiation, and used as a cell-based therapy for genetic
20 diseases. This technology is creating enormous interest because the nature of both the genetic modification and the transplanted cell type would be unrestricted. This circumvents important limitations to current gene therapy protocols. A barrier to this technology is the requirement for transplanted cells to evade immune rejection. One solution to this might be creation of a generic
25 human 'donor' ES cell line, modified to escape immune surveillance. A preferred approach is the development of techniques, which allow the isolation of 'individual-specific' pluripotent cells. This can potentially be achieved via the use of nuclear transfer technology where pluripotent cells are isolated from viable embryos generated by injecting somatic cell karyoplasts into an oocyte
30 cytoplasm. This approach raises important ethical considerations. An alternative



5

10

20

25

of pluripotent cells, partially differentiated cells and differentiated cells have been limited. Nuclear transfer has been one of the approaches suggested in the prior art to achieve reprogramming of differentiated somatic cells. However nuclear transfer approaches have been inefficient due at least in part, to
5 difficulties in reprogramming. Furthermore there are major ethical problems with the use of human oocytes in reprogramming human somatic cells.

There have also been problems in controlling the differentiation of pluripotent cells along defined differentiation pathways.

There have also been difficulties associated with primary somatic cells
10 in vitro. In particular it has been difficult to maintain primary cells in culture for prolonged periods, due to biological mechanisms that limit the number of proliferation rounds that such cells can undergo. The consequence of this limitation, termed the Hayflich limit, is that the ability to genetically modify primary or untransformed cells in vitro has been restricted. This limitation has
15 restricted the utility of these cells for commercial, agricultural or medical benefit.

It is accordingly an object of the present invention to overcome or at least alleviate one or more of the difficulties or deficiencies associated with the prior art.

20 Applicants have discovered that the unusual mode of cell cycle regulation seen in pluripotent cells is an intrinsic feature of the pluripotent state and that maintenance of this state is dependent, in part at least, on the behaviour and properties of molecules that are involved in cell cycle regulation/cell proliferation. Accordingly, these molecules may be used to
25 identify, isolate and maintain populations of pluripotent cells in vitro, that are otherwise unmanageable. This may also facilitate the identification and isolation of cells that have dedifferentiated back to a less differentiated state.

Identification of pluripotent cells and multipotent cells

Accordingly, in a first aspect of the present invention there is provided a method for identifying pluripotent cells on the basis of their cell cycle dynamics and expression and/or activity of cell cycle regulatory molecules.

5 Specifically, there is provided a method for identifying pluripotent or pluripotent-related cells which method includes

analysing the cell population for pluripotent cell cycle characteristics including one or more of

- pluripotent-specific cell cycle structure;
- 10 pluripotent-specific expression and/or activity of cell cycle regulatory molecules; and
- phosphorylation status of a tumour-suppressor protein(s).

In a preferred aspect the identification method may include measuring expression of cell cycle regulatory molecules, including a cyclin(s), a cyclin-
15 dependent protein kinase (Cdk), a Cdk inhibitor, upstream regulators of said molecules or biochemical targets thereof. The method may also include determining the phosphorylation state of a tumour suppressor protein such as pRb and other family members such as p107 and p130.

20 It was known in the prior art that ES cells proliferate rapidly, with a cell cycle of about 10-12 hours in vitro. However little was known previously about the cell cycle structure, and the molecular events regulating the cell cycle in pluripotent cells.

The applicant has surprisingly found that ES cells and other pluripotent cells including EPL cells, and pluripotent cells in the developing embryo have a
25 different cell cycle structure compared to differentiated cells. Pluripotent cells spend the majority of time (~65%) in S phase, and short proportions of the cell cycle in G1 (~15%) and G2/M (~20%) phases. In the case of G2 and M-

Applicant has also found that the tumour suppressor Retinoblastoma protein (pRb) is maintained in an inactive state by phosphorylation in

- elevated constitutive expression and/or activity of cyclin A;
- substantial reduction in, or absence of, expression of Cdk inhibitor p16,
- substantial reduction in, or absence of, expression of Cdk inhibitors p21
and/or p27; and
- 5 presence of an inactive (phosphorylated) tumour suppressor protein.

Cells with properties similar to the cell cycle characteristics found in pluripotent cells are identified as pluripotent cells.

Preferably the pluripotent or pluripotent-related cells are identified by the presence of one or more of the following characteristics:

- 10 a pluripotent-specific cell cycle exhibiting a rapid cycle with short gap phases;
- elevated constitutive expression and/or activity of cyclin E;
- elevated constitutive expression and/or activity of cyclin A;
- pluripotent-specific expression of Cdk inhibitors; and
- 15 presence of a phosphorylated tumour suppressor protein.

In a further embodiment of this aspect of the present invention, the pluripotent-related cells may include multipotent cells (such as haemopoietic stem cells and neural stem cells). The multipotent cells may be derived by partial differentiation of pluripotent cells and which are capable of
20 differentiating further into a number of different cell types, which may have all or some of the above cell cycle features.

In a further aspect of the present invention, there is provided
a method of identifying differentiating cells, which method includes
analysing the cell population for differentiating cell cycle characteristics
25 including

- 16 -

differentiation-specific cell cycle structure;
differentiation-specific expression and/or
activity of cell cycle regulatory molecules, including
the presence of active tumour suppressor proteins.

- 5 In one embodiment, the identification method may be utilised to monitor the onset of differentiation.

The cell cycle structure, expression and activity of cell cycle regulatory molecules are altered significantly during differentiation. The alterations may include one or more of the following:

- 10 In relation to the cell cycle, the gap phases of the cell cycle become more prominent, and the rapidity of the cell cycle slows.

Differentiating cells are increasingly dependent on tumour suppressor proteins, such as pRb and other family members for regulation of the cell cycle.

- 15 cyclin E-is significantly downregulated in differentiating cells, through reduced gene expression, and reduced protein stability.

cyclin A-associated activities are also reduced.

Cdk 2 activities become cell cycle regulated.

- 20 The activities of other cell cycle regulatory molecules, such as the INK Cdk inhibitor p16 and/or other INK family members are upregulated.

The activities of other Cdk inhibitors p21 and/or p27 are upregulated.

Accordingly differentiating cells may be identified by the presence of

In one embodiment, maintenance and proliferation of pluripotent cells in

25 The lifespan of pluripotent cells *in vitro* may also be prolonged by manipulating the activities and expression of tumour suppressor molecules, such as pRb, and related activities, p107 and p130. In general cell cycle entry may be promoted by hyperphosphorylation and inactivation of tumour suppressor proteins such as pRb, leading to activation of the E2F family of transcription factors. Inactivation of pRb may be achieved by

hyperphosphorylation, or by other approaches that include antisense technology, or gene inactivation. Similarly prolonged lifespan, proliferation and continued maintenance of differentiation status may be achieved by constitutive expression or activity of E2F transcription factors. For example,
5 constitutive E2F activity would occur when E2F is manipulated so that it no longer interacts with nor inhibited by pRb or other tumour suppressors.

The pluripotent cells may be of any suitable type and may be *in vitro* or *in vivo*. Preferably, the pluripotent cells are selected from one or more of the group consisting of epiblast cells, ES cells, EPL cells (as described in
10 International Patent application PCT/AU99/0265), primordial germ cells (PGCs), or embryonic carcinoma (EC) cells.

Similarly the multipotent cells may be of any suitable type and may be *in vitro* or *in vivo*. They may be any partially differentiated cell type, including such cells as haematopoietic stem cells and neural stem cells.

15 Where the cell cycle regulatory molecule includes a cyclin, the cyclin may be of any suitable type. Preferably the cyclin is cyclin D, cyclin E, cyclin A or a molecule exhibiting similar activity (e.g. virally encoded cyclins that are not inhibited by Cdk inhibitors), or a functionally active fragment or analogue thereof. The cyclin-dependent protein kinase may be of any suitable type and
20 includes biochemical activities with similar properties. Preferably the Cdk is Cdk4, Cdk6, or Cdk2, or a molecule exhibiting similar activity, or a functionally active fragment or analogue thereof.

The upstream regulatory pathways may be components of signalling pathways that, in some cases, are known to modulate aspects of cell
25 behaviour such as, but not limited to, cell proliferation, the cell cycle and differentiation status. This would include and is not limited to molecules such as myc family members, including c-myc, l-myc and n-myc, Ras, Raf, MAP kinase, Rho and other signalling pathways. Ultimately the activity of these

molecules would serve to provide and maintain intracellular activities required for stem cell maintenance separate to or, in conjunction with, the cell cycle functions described.

5 The activities of said cell cycle regulatory molecules may be manipulated by any suitable technique. Such techniques include but are not limited to manipulation of expression of said proteins, including manipulation of gene expression for example by transformation with expression constructs, antisense technology or fusion protein technology where the fusion protein includes a transduction domain linked to the cell cycle regulatory molecule, or
10 any other protein delivery system such as electroporation or lipofection. Such techniques are well known to those skilled in the art and are described in, for example, Sambrook et al (1989), the entire disclosure of which is incorporated herein by reference.

In a preferred form the manipulated cell cycle regulatory molecule is an
15 upstream regulator of cyclin E and/or cyclin A, such as c-myc and other members of the myc family. In this preferred form c-myc activity is upregulated by ectopic gene expression, or most preferably by protein transduction, using a fusion protein that includes a transduction domain linked to the cell cycle regulatory molecule, or any other protein delivery system such as
20 electroporation or lipofection.

Methods for extending the lifespan in vitro, of primary cells and untransformed cells

There have also been difficulties associated with the short lifespan of primary somatic cells and untransformed cells in vitro. In particular there are
25 biological mechanisms that limit the number of proliferation rounds that such cells can undergo. This difficulty is termed the Hayflich limit. One consequence of this difficulty is that the ability to genetically modify primary or untransformed cells in vitro is restricted, limiting their potential applications in a range of technologies, including nuclear transfer.

The lifespan *in vitro* of primary and untransformed cells may be prolonged by manipulation of cell cycle regulatory activities.

Accordingly, in accordance with this aspect of the present invention, there is provided a method of regulating the cell cycle of primary and
5 untransformed cells, which method includes

manipulating the expression and/or activity of a cell cycle regulatory molecule, such that the proliferation and maintenance of differentiated cells are reduced, wherein the regulatory molecule is selected from one or more of the groups consisting of a cyclin, a cyclin-dependent protein kinase (Cdk), a Cdk
10 inhibitor, upstream regulators thereof or biochemical targets thereof and/or tumour suppressor protein, and molecules displaying similar activities, in a primary or untransformed cell.

In an alternative or supplementary embodiment, the lifespan of such cells *in vitro* may be prolonged by manipulating the activities and expression of
15 tumour suppressor molecules, such as pRb, and related activities, p107 and p130. In general, cell cycle entry may be promoted by hyperphosphorylation and inactivation of tumour suppressor proteins such as pRb, leading to activation of the E2F family of transcription factors. Inactivation of pRb may be achieved by hyperphosphorylation, or by other approaches that include
20 antisense technology, or gene inactivation. Similarly prolonged lifespan, proliferation and continued maintenance may be achieved by constitutive expression or activity of E2F transcription factors. For example, constitutive E2F activity would occur when E2F is manipulated so that it no longer interacts with nor inhibited by pRb or other tumour suppressors.

25 ***Methods for selecting pluripotent cells***

In another aspect of the present invention there is provided a method of selecting for pluripotent cells from a mixed cell population including pluripotent cells, and differentiated cells which method includes

manipulating the activity of a cell cycle regulatory molecule, such that the proliferation and maintenance of differentiated cells are reduced, wherein the regulatory molecule is selected from one or more of the groups consisting of a cyclin, a cyclin-dependent protein kinase (Cdk), a Cdk inhibitor, upstream
 5 regulators thereof or biochemical targets thereof and/or tumour suppressor protein, and molecules displaying similar activities, in a pluripotent or multipotent cell.

The differentiated cells may be already present in the cell population, or derived by differentiation of pluripotent cells. Elimination of differentiated cells
 10 from the population would in itself improve overall pluripotent cell stability, since differentiated cells are capable of promoting differentiation of pluripotent cells. Hence this method confers an additional advantage in enhancing pluripotent cell stability.

The preferential inhibition of differentiated cell maintenance and
 15 proliferation without inhibiting pluripotent cell maintenance and proliferation may be achieved by manipulation of any of the cell cycle regulatory molecules including cyclins, cyclin-dependent protein kinases, and Cdk inhibitors, or their upstream regulators or down stream targets. Upstream regulators include the protooncogenes from the *myc* family, which upregulate cyclin E-associated
 20 activities and cyclin A-associated activity. Other upstream signalling pathways such as Ras, Raf, MAP kinase, or Rho may also be used.

In a preferred embodiment preferential inhibition of differentiated cell maintenance and proliferation is achieved by manipulating Cdk inhibitor activity. Pluripotent cells would remain insensitive to the manipulated Cdk
 25 inhibitor activity, whereas progress of differentiated cells through the cell cycle is preferentially inhibited.

In a particularly preferred embodiment, the activity of the Cdk inhibitor p16 is manipulated. Constitutive p16 activity in pluripotent cells does not inhibit

In yet another embodiment, preferential inhibition of differentiated cell maintenance and proliferation could be used in combination with approaches

to promote pluripotent cell maintenance and proliferation. For example a Cdk inhibitor such as p16 may be used to inhibit differentiated cell maintenance and proliferation, in combination with cyclin E and/or cyclin A, or upstream regulators of cyclin E- and/or cyclin A-associated activities such as the myc
5 family of proto-oncogenes to promote maintenance and proliferation of those pluripotent cells selected on the basis of their insensitivity to p16 or other INK family members with similar activity.

In another form, manipulation of cell cycle regulatory molecules can be used to preferentially select for partially differentiated cells from a mixed cell
10 population comprised of partially differentiated cells and differentiated cells.

Use of this approach to trap cells that spontaneously revert to a less differentiated state.

More particularly, these methods can be exploited to obtain cells that have reverted from a differentiated state to a less differentiated state, in a
15 manner that is independent of nuclear transfer. Reversion to a less differentiated state may occur spontaneously. In differentiated cell populations, including multipotent cell populations, it is probable that spontaneous reversion to a less differentiated state occurs at a low frequency, or rarely. Reverted cells may be trapped in their less differentiated state, and
20 selected for by the previously described methods.

In one embodiment enforced Cdk inhibitor expression, and in particular p16 expression (p16 or other INK family members with similar activity is the preferred Cdk inhibitor) is achieved in a differentiated cell population by techniques that include ectopic gene expression or any other means. Most
25 preferably Cdk inhibitor activity is achieved by protein transduction, using a fusion protein that includes a transduction domain linked to p16, or any other protein delivery system such as electroporation or lipofection. Differentiated cells, which rely on cyclin D-associated activities for proliferation, are sensitive to p16 activity, and fail to proliferate, and undergo cell death. Rare individual

Non spontaneous reversion may be initiated by placing multipotent or differentiated cells in an environment in vitro that promotes programming. The methods described in this application for the selection of pluripotent cells or multipotent cells can be adapted to trap and temporarily lock cells in the transitional stages of reprogramming. One likely possibility is that environmental signals are able to direct reprogramming through a transitional pluripotent state, and subsequent differentiation through all germ line lineages. In this approach the signals that promote reprogramming are derived from pluripotent cells, or from cells in early differentiation, as occurs in embryoid

- 27 -

The Cdk inhibitors may also be used in combination with other methods for maintenance and proliferation of pluripotent cells, such as manipulation of cyclin E, or upstream regulators such as myc family etc.

5 These methods may also be used to capture and maintain cells in a transitional multipotent state, formed by reprogramming in response to an inductive environment.

Methods for reprogramming of differentiated or partially differentiated cells, to a less differentiated state, including to a state of pluripotency

10 Manipulation of the activities of cell cycle regulatory molecules also provides approaches for the dedifferentiation of differentiated cells.

Accordingly in this aspect of the present invention, there is provided a method for reprogramming of differentiated or partially differentiated cells to a less differentiated state which method includes

15 manipulating the expression and/or activity of a cell cycle regulatory molecule, including a regulatory molecule selected from one or more of the groups consisting of a cyclin, a cyclin-dependent protein kinase (Cdk), a Cdk inhibitor, upstream regulators thereof or biochemical targets thereof and/or tumour suppressor protein, and molecules displaying similar activities, in said
20 differentiated or partially differentiated cells.

For example, cell cycle regulatory molecules or their upstream regulators or downstream targets are manipulated in differentiated cells, so that their cell cycle properties assume at least some of the cell cycle properties of pluripotent or multipotent cells.

25 Dedifferentiation may be achieved by manipulation of any of the cell cycle regulatory molecules including cyclins, cyclin-dependent protein kinases,

and Cdk inhibitors, or their upstream regulators or down stream targets. Upstream regulators include the protooncogenes from the myc family, which upregulate cyclin E-associated activities. Other upstream signalling pathways such as Ras, Raf, MAP kinase, or Rho may also be used.

- 5 Examples include cyclin(s), particularly upregulation of cyclin E, cyclin A; cyclin-dependent protein kinases, particularly Cdk2; Cdk inhibitors; and upstream regulators of cyclin E- and/or cyclin A-associated activities, and other upstream pathways; particularly myc.

- 10 Dedifferentiation may be achieved by manipulation alone, or in combination with other procedures that promote dedifferentiation, such as nuclear transfer.

- 15 In the prior art, the efficiency of nuclear transfer is low, and only a small proportion of nuclear transfer embryos develop to live animals. This inefficiency is due largely to problems with reprogramming of the somatic nucleus to a pluripotent state after transfer to the recipient oocyte. Reprogramming efficiency can be improved by manipulating cell cycle properties of donor somatic cells so that they more closely resemble cell cycle features of a pluripotent cell, prior to nuclear transfer.

- 20 In a preferred embodiment, manipulation of cell cycle regulation in differentiated cells or partially differentiated cells in this way, may prime cells so that they are more readily dedifferentiated by nuclear transfer. Pluripotent cells may be formed from manipulated cells by fusion with enucleated oocytes, or cytoplasts derived from pluripotent cells.

- 25 ***Methods for regulating differentiation of pluripotent cells by manipulation of cell cycle.***

the activity of Cdk2 becomes cell cycle regulated; or
upregulating the activities and/or expression of CdK inhibitors.

Methods include:

5 increasing the role of tumour suppressor proteins, such as pRb and
other family members in regulation of the cell cycle.

cyclin E-downregulation by methods that include reduced gene
expression (eg antisense technology) and/or reduced protein stability.

10 reducing cyclin A-associated activities, e.g. by methods that include
reduced gene expression (eg antisense technology) and/or reduced protein
stability.

Cdk 2 activities become cell cycle regulated.

Upregulate the expression and/or activities of Cdk inhibitors such as
p16, p21 and p27.

15 The present invention will now be more fully described with reference to
the accompanying figures and examples. It should be understood, however,
that the description following is illustrative only and should not be taken in any
way as a restriction on the generality of the invention described above.

In the Figures:

Figure 1 illustrates regulation of the cell cycle.

20 Figure 2 illustrates remodelling of the cell cycle during embryonic
development.

Figure 3 illustrates flow cytometry analysis of pluripotent embryonic
epiblast cells isolated from day 6.5dpc embryos.

Figure 4 illustrates comparison of cell cycle profiles between 6.5dpc embryonic epiblast, ES, EPL and NIH 3T3 fibroblasts.

The general cell cycle structure and cell cycle length are indicated. Note the relatively short cell cycle length and lack of fully formed gap phases in
5 pluripotent ES, EPL and epiblast cells.

Figure 5 illustrates cell cycle remodeling during differentiation of cells in embryoid bodies.

ES cells or EPL cells grown as EPL embryoid bodies in the absence of LIF were fixed and stained with propidium iodide. EPL bodies grown in the
10 absence of LIF were harvested to evaluate changes in cell cycle structure associated with differentiation. Note the relative increase in the proportion of G1 cells and a decrease in the proportion of S-phase cells as cells differentiate.

Figure 6 illustrates E2F target genes are not cell cycle regulated in ES
15 cells.

Top panel: ES cells were synchronized by the nocodazole-aphidicolin block –release protocol. Following release from the G1/S aphidicolin block, cells were harvested, RNA was prepared and resolved on 1% formaldehyde agarose gels, blotted onto a nitrocellulose membrane and probed with a P³²-
20 labelled cyclinE, RRMP-2 or mGAP cDNA fragment. The timing of entry into S-phase for synchronous ES cell populations is indicated. Lower panel:

E2F4. The major pRb-like activity is shown to be p107, by using the super-shift assay, with an anti-p107 antibody.

Figure 9 illustrates Cyclin E and cyclin B associated activities in ES cells and NIH3T3 cells.

5 Whole cell extracts from ES or NIH 3T3 cells were prepared from unsynchronized (untreated), +aphidicolin treated (G1/S block) or nocodazole blocked (G2/M) cells. 50• g total protein from each extract was used in immunoprecipitation assays with anti-cyclinE or anti-cyclinB antibody. Kinase activities in the immunoprecipitates were evaluated using recombinant GST-
10 pRb or purified histone H1 as an in vitro substrate.

Figure 10 illustrates levels of cyclin E and cyclin A are not cell cycle regulated in ES cells.

ES cells were synchronized and released into S-phase. At 90 minute intervals, samples of cells were collected, whole cell extracts were prepared
15 and levels of Cdk4, cyclinE, cyclinB, cyclinA, Cdk2 and Cdc2 detected by Western blotting. Note that in ES cells, cyclinB levels vary in a cell-cycle dependent manner (ie levels peak late in G2/M-phase). CyclinE and cyclin A levels do not vary throughout the cell cycle however.

Figure 11 illustrates Cyclin E and cyclin A activities and mRNA
20 pluripotent cells and during differentiation in embryoid bodies.

were stained for alkaline phosphatase activity. The data shows the % colonies staining positive for alkaline phosphatase. Each data point is the average of assays performed in duplicate.

Figure 14 illustrates ectopic expression of c-myc severely reduces the
5 ability of pluripotent cells to differentiate.

ES cells were grown into EPL embryoid bodies and their ability to differentiate monitored using molecular markers (Oct4, pluripotency; brachyury, nascent mesoderm). RNA was prepared from embryoid bodies at different times after withdrawal of LIF (1-5 days). RNA was subject to agarose
10 gel electrophoresis and individual RNAs detected by Northern blot. Note that c-myc expressing cells do not switch off Oct4 (and hence, retain pluripotency) and only turn on the differentiation marker (brachyury) at low levels. Data for two c-myc and one control cell line is shown. Levels of ribosomal RNA are shown in the right hand panel.

15 Figure 15 illustrates Tat-p16 fusion protein selectively inhibits the growth of NIH3T3 fibroblasts but not ES cells.

Varying concentrations of TAT-p16 or TAT-p16mut (0-150• g/ml) were added to sub-confluent cultures of NIH3T3 or ES cells. Cells were counted at time points after addition of TAT fusion protein. Each experiment was
20 performed in duplicate and data points represent average values.

EXAMPLE 1: The cell cycle profile of pluripotent cells *in vitro* and

epiblast cells *in vivo* are similar, and are strikingly different to the profile in differentiated cells.

To demonstrate that the mode of cell cycle regulation is fundamentally different in pluripotent and differentiated cells, we performed flow cytometry analysis (based on DNA content) on purified populations of epiblast cells derived from mouse embryos and on pluripotent cells derived from *in vitro* culture.

Materials and Methods

Isolation of pluripotent cells from 6.5dpc embryos:

Isolation of pluripotent cells from 6.5dpc embryos were by dissection from time-mated Swiss mice, Reichardt's membrane and the extra-embryonic part removed, before the visceral endoderm peeled away from the epiblast by pipetting the embryo up and down with a narrow bore pipette. Each epiblast at this stage of development consists of between 750-1000 cells making it necessary to pool approximately 10 embryos to generate sufficient cell numbers for flow cytometry analysis. After generating a single cell suspension, cells were then fixed and stained with propidium iodide (described in Dunphy 1997) for flow cytometry analysis on the basis of cellular DNA content (Figure 3). This analysis allows the relative proportion of time spent in G1, S-phase and G2/M to be determined.

Cell culture *in vitro*:

Mouse embryonic stem (ES) cells were cultured in the absence of feeders on tissue-culture grade plastic-ware (Falcon) pre-treated with 0.2% gelatin-PBS for a minimum of 30 minutes. Cells were cultured in Dulbecco's Modified Eagles Medium (GIBCO BRL) pH7.4 containing high glucose. To
5 generate the final DMEM used for cell culture this was supplemented with 10% foetal calf serum (FCS; Commonwealth Serum Laboratories), 40mg/ml gentamycin, 1 mM L-glutamine and 0.1 mM β -mercaptoethanol (β -ME). DMEM was supplemented with 1000 units of LIF under 10% CO₂ in a humidified incubator where appropriate. Routine tissue culture was performed
10 as described by Smith (1991).

LIF was routinely produced from COS-1 (ATCC CRL-1650) cells transfected with a mouse LIF expression vector plasmid, pDR10, as described by Smith (1991). COS-1 cells were transfected by electroporation using a Bio Rad Gene Pulsar at 270 Volts and a capacitance of 250 μ FD. Transfected
15 cells were plated at 7×10^4 cells/cm² in DMEM, pH 7.4, containing high glucose and supplemented with 10% FCS, 40 μ g/ml gentamycin and 1mM L-glutamine. Medium was collected and assayed for LIF expression as described by Smith (1991). Alternatively, medium was supplemented with 1000 units of recombinant LIF (ESGRO, AMRAD).

20 E14, CGR8, E14TG2a ES cells (Hooper et al 1987), MBL5 (Pease et al 1990) and D3 (Doetschman et al., 1985) ES cells were cultured in DMEM supplemented with LIF as described above. Embryonal carcinoma cell lines were F9 and P19 (ATCC CRL 1825). Mouse embryo fibroblasts were isolated

from day 12.5 Swiss mouse embryos as described in Hogan et al 1994. Other cell lines including NIH3T3 fibroblasts (ATCC CRL1658), Balb/c 3T3 clone A31 (ATCC CCL 163) were grown in DMEM supplemented with 10% FCS.

Hep G2 cells (Knowles et al., 1980; ATCC HB-8065) were maintained in culture in DMEM and passaged at confluence. To condition medium (MEDII) Hep G2 cells were seeded into DMEM at a density of 5×10^4 cells/cm². Medium was collected after 4-5 days, sterilized by filtration through a 0.22 μ m membrane and supplemented with 0.1 mM β -ME before use. MEDII was stored at 4°C for 1-2 weeks or at -20°C for up to 6 months with apparent loss of activity.

EPL cells were formed and maintained in media containing 50% MEDII conditioned medium in DMEM with or without the addition of LIF. EPL formation was apparent with the addition of between 10 and 80% MEDII, with optimal culture conditions at 50% MEDII.

Pluripotent EPL cells were formed from ES cells and maintained as follows;

Adherent cultures: ES cells were seeded at a density of 1×10^5 cells/cm² onto tissue culture grade plastic-ware (Falcon) pre-treated with 0.2% gelatin/PBS for a minimum of 30 minutes in DMEM containing 50% MEDII as described above. EPL cells were maintained in 50% MEDII using routine tissue culture techniques (as described by Smith, 1991).

Suspension aggregates: ES cells were seeded at a density of 1×10^5 cells/cm² in suspension culture in bacterial petri dishes in DMEM containing 50% MEDII as described above. The resulting EPL cell aggregates were split 1:2 after 2 days and seeded into fresh DMEM containing 50% MEDII. Growth of embryoid bodies (EPL or ES) were described in Rathjen et al 1999.

Flow cytometry analysis: Cell cycles profiles were determined by propidium iodide staining of fixed cells as described in Dunphy (1997).

Western blot analysis: Antibodies used for Western blot and immunoprecipitation analysis were as follows. cyclin E (Santa Cruz sc-481 or sc-198), p16 (Santa Cruz sc-1207), pRb (Becton Dickinson 14001A), p107 (Santa Cruz sc-318), E2F-4 (Santa Cruz sc-866), Cdk2 (Santa Cruz sc-163), p27 (Santa Cruz sc-528), p21 (Santa Cruz sc-397). HRP-conjugated secondary antibodies raised against total rabbit or mouse immunoglobulins were from Dako Corporation (P217 and P0260, respectively). Detection of proteins in Western blot analysis was by the Pierce ECL kit (#34080).

Results

Our results reveal some striking features of epiblast, ES and EPL cell cycles relative to cell cycles of differentiated somatic cells. First, it is apparent that pluripotent cells, including those of the epiblast, spend the majority of their time in S-phase (~65%) and short proportions of the cell cycle in G1 (~15%) and G2/M (~20%) phases (Figures 3,4). In the case of G2 and M-phase cells

The potential consequences of shortened G1 and G2/M phases in pluripotent cells, could be decreased cell cycle length and loss of regulatory controls such as checkpoints. Molecules known to modulate the length of the gap phases and of cell cycle length when ectopically expressed include cyclin E (Resnitzky et al 1994) and c-myc (Kam et al 1989), respectively. The

- 41 -

molecular mechanisms underlying changes in cell cycle structure and rates of cell division in pluripotent cells and multipotent cells have not been addressed previously and neither has the relevance of this to the establishment and maintenance of the pluripotent state.

5 **EXAMPLE 2: Regulation of the cell cycle in pluripotent cells.**

Material and Methods

All cells and tissue culture techniques were as described in Example 1 unless otherwise stated.

Northern Blot Analysis and in vitro Cdk assays:

10 Cytoplasmic RNA preparation and Northern blot analysis were by standard methods (Sambrook et al., 1989). In vitro kinase assays were performed as described in Dunphy (1997).

Synchronization of ES cells

15 A synchronization procedure, taking into account the unusual nature of the pluripotent cell cycle structure, was developed and optimized. ES cells were first plated as single cell suspensions at 5×10^5 cells/ml onto gelatin-coated petri dishes (see Example 1). After 12-14 hours culture in DMEM plus LIF, the microtubule depolymerizing agent nocodazole (Sigma), was added to a final concentration of 45ng/ml. After 8 hours, cells were washed three times

- 42 -

in DMEM (37°C) and then incubated for a further 5-8 hours in DMEM plus LIF supplemented with 5• g/ml aphidicolin (Sigma), which blocked the cell population at the G1/S border. Cells were released from the G1/S cell cycle block by washing three times in DMEM supplemented with LIF (warmed to 5 37°C). During each wash cycle, cells were kept under standard culture conditions for five minutes between media changes. Cell cycle arrest, synchrony and cell cycle transit was evaluated by flow cytometry (see Example 1) to evaluate cell DNA content (data not shown). After release from cell cycle blocks, cell samples were taken and used for cell cycle analysis (for example; 10 protein extracts, RNA extractions, flow cytometry analysis).

Results

pRb activity in pluripotent cell cycles

The retinoblastoma tumour suppressor protein, pRb, and its family members (p107, p130) are key regulators of the G1-S transition and are 15 crucial for processes that normally control cell proliferation. Their activity is controlled by the activity of cell cycle regulated Cdk activities which impose a distinct pattern of cell cycle regulated phosphorylation on pRb-family members (Dyson 1998). Underphosphorylated pRb is generally accepted to be the active form, capable of interacting with cellular targets such as the E2F 20 family of transcription factors (Dyson 1998). The hyperphosphorylated form is the inactive species, and one measure of this is the loss of pRb's ability to associate with E2F transcription factors. Loss of pRb function in tumour cells is commonly associated with deregulated E2F activity and has several

20 The ability of pRb to inhibit E2F transcription factors was determined by measuring the expression of E2F-target genes that are normally activated during the G1-S period of the cell cycle (Figure 6). Northern analysis was conducted on RNA isolated from synchronised ES cells, and probed for cyclin

- 44 -

E and RRMP-2 mRNA (RRMP-2 refers to a ribonucleotide reductase subunit).

Both these genes are normally transcribed by E2F transcription factors and are cell cycle regulated in differentiated somatic cells. mGAP mRNA levels remain constant throughout the cell cycle, and was used as a control to show
5 relative RNA loading. Results shown in Figure 6 indicate that these genes are active throughout the cell cycle and not subject to any cell cycle-regulated repression/activation. Hence, E2F target genes are constitutively active, and pRb (and p107, p130) regulation of E2F activity does not exist in ES cells. This contrasts the general situation in primary and established cell lines other than
10 pluripotent cells, where E2F target genes are tightly cell cycle-regulated through Cdk regulation of pRb-family proteins. Hence, this shows a fundamental difference in the cell cycle regulatory pathways that operate in pluripotent stem cells.

The ability of pRb to interact with E2F transcription factors was tested
15 using gel mobility shift analysis (Figure 7). Our work identifies E2F-4 as the major E2F DNA binding activity in ES cell nuclear extracts; it is this E2F activity that is shifted by an anti-E2F antibody in gel shift mobility assays (Figure 8). Remarkably, the vast majority of E2F activity in pluripotent cells was in the 'free' state, uncomplexed with pRb family proteins (Figure 7). No E2F-pRb
20 complexes were detected by band shift analysis using supershifting anti-pRb antibodies to probe the composition of protein-DNA complexes (data not shown). Instead, a small fraction of E2F-4 is associated with p107, (another member of the pRb tumour suppressor family) although the majority of E2F-4

To learn more about proliferative control in pluripotent cells, we have

investigated Cdk activities associated with Cdk2. The activity of Cdk2-cyclinE activity was evaluated in asynchronous (untreated), G2/M (nocodazole) or G1/S (aphidicolin) blocked cells. This experiment demonstrates an unusual mode of Cdk regulation in pluripotent ES and EPL cells because cyclin E-Cdk2

5 is active at unusually high levels (see direct comparison between pluripotent cells and early passage MEFs, with equal protein input in Figure 11) and moreover, is promiscuously active throughout the cell cycle (Figure 9). Under normal circumstances cyclin E-associated Cdk activities are tightly cell cycle regulated, peaking in G1/early S-phase. In pluripotent cells however, cyclin E-

10 kinase activity is still active in a nocodazole arrest (G2/M block) which is normally associated with inactive cyclin E-kinases in differentiated cells. The mitotic cyclin B-associated kinase activity behaves as expected however, and is more active in G2/M cells than at other points in the cell cycle. This is consistent with the situation in differentiated cells (such as NIH 3T3

15 fibroblasts). Hence, the pluripotent state is associated with an unusual pattern of activity for some Cdk activities (cyclin E, cyclin A/Cdk2), but not all (Cdk1-cyclin B). Similar results were obtained in F9 and P19 pluripotent embryonal carcinoma cells and pluripotent EPL cells (data not shown), indicating that this unusual mode of Cdk regulation is a general property of pluripotent cells.

20 Levels of cyclin E-Cdk2 pRb kinase activity are of comparable activity in G2/M cells and G1 cells and are significantly higher in magnitude than that found in MEFs, NIH 3T3s and other cell lines tested (Figure 9 and data not shown). After adjustment of total protein input, the amount of Cdk2 activity in pluripotent ES, EC and EPL cells was determined to be 20-50X higher than in

low passage mouse embryo fibroblasts (data not shown). The overall characteristics of Cdk2 regulation in pluripotent cells deviates significantly from the normal pattern of cyclin E-Cdk2 activity in differentiated somatic cells, where cyclin E-Cdk activity increases during G1 and collapses during S-phase, as a result of tight cell cycle regulatory mechanisms. Promiscuous Cdk2-cyclin A activity was also seen- ie absence of cell cycle regulation and elevated levels relative to primary and immortalized cell lines (see Figure 11 and data not shown).

In support of the continued cyclin E-Cdk2 activity throughout the cell cycle, cyclin E protein was shown to be present throughout the cell cycle in synchronised cells by Western analysis (Figure 10), at elevated levels. This contrasts the situation in other cell types where cyclin E levels are strictly cell cycle regulated due to transcriptional and post-transcriptional mechanisms (Sherr 1994). Mitotic cyclins however, such as cyclin B, exhibit cell-cycle dependency (Figure 10) as is typically seen in differentiated, non-pluripotent cells. This lack of cell cycle regulated control of cyclin E and cyclin A at the protein level is multifactorial and can be explained in part at least by, (i) the lack of cell cycle regulation of the cyclin E/cyclin A transcription (the cyclin E and cyclin A genes are E2F target genes) and, (ii) enhanced stability of the cyclin E/cyclin A proteins in pluripotent cells (data not shown).

In the early embryo, our data derived by immunohistochemistry shows that prior to gastrulation, cyclin E staining is intense in all cells of the epiblast (data not shown). This is consistent with the observation in ES cells that cyclin

E levels are invariant throughout the cell cycle. This contrasts the staining pattern in the surrounding visceral endoderm layer (data not shown) where general staining is less intense and only detected in a fraction of cells (this is probably coincident with G1 and S-phase cells). The precocious activities of Cdk2-cyclin E and Cdk2-cyclin A most likely underpin the constitutively phosphorylated state of pRb in pluripotent stem cells, hence explaining pRbs biochemical inactivity and constitutive activity of E2F target genes.

Additional work has shown that precocious cyclin E-Cdk2 activity is a special feature of the pluripotent state. As pluripotent ES and EPL cells are allowed to differentiate (following withdrawal of LIF), cyclin E-Cdk2 activity was monitored in relation to well established differentiation markers for pluripotency, nascent mesoderm and primitive ectoderm: Oct4, brachyury and FGF-5, respectively (Figure 11). Our data shows that as pluripotent cells are allowed to differentiate (withdrawal of LIF), several key events occur. (1) cell cycle structure gets remodelled such that the gap phases become longer (see Figure 5), (2) cell cycle length becomes longer as a consequence of (1), (3) Cdk2 activities become severely down-regulated as cells lose pluripotency (Figure 11), (4) this is associated with down-regulation of cyclin E protein levels (Cdk2 protein remains constant, Figure 11) and (5) as part of the differentiation program, Cdk2-cyclin E and Cdk2-cyclin A activities become cell cycle regulated (data not shown). Loss of precocious cyclin E-Cdk2 and cyclin A-Cdk2 activities and establishment of their cell cycle regulation are mechanistically linked to the loss of pluripotency (see Figure 11).

The apparent unrestrained activity of Cdk2 kinases that persists in pluripotent cells throughout the cell cycle, occurs in the absence of two Cdk inhibitory molecules, p21 and p27 (Figure 12), known to modulate Cdk2 activity (Sherr and Roberts 1995). Besides being Cdk inhibitors for Cdk2
5 kinase complexes, p21 and p27 are established assembly factors for the Cdk4,6-cyclin D complex (Cheng et al 1999). These latter complexes are principally involved in control of the R-point by phosphorylating pRb and it's family members. In the absence of p21 and p27, little or no active Cdk4-cyclin D assembly activity would be expected, implying that these activities do not
10 play a major role in the control of pluripotent cell proliferation. Upon cellular differentiation (after loss of pluripotency, for example) however, when p21 and p27 first become present, it would be expected that Cdk4,6-cyclin D activities would become crucial. Hence the requirement for Cdk4,6-cyclinD activities occurs as cells lose pluripotency and as Cdk2 activities become cell cycle
15 regulated.

The presence of pRb in a biochemically inactive, hyper-phosphorylated state, suggests from other prescedents that pluripotent cells lack a functional R-point. Because these cells lack functional pRb, they should exhibit a reduced requirement for Cdk4,6 kinase activities (see Jiang et al
20 1998 and references therein). This situation can be satisfied if Cdk2 activities are elevated, as in the case of some tumour cell lines. Our data indicate that pluripotent cells also will also have a reduced requirement for Cdk4,6 activities as they also have elevated Cdk2 activities and lack functional pRb. The

absence of pRb activity is likely to be through a mechanism where precocious Cdk2-cyclin E/A holds it in a constitutively phosphorylated state, independent of Cdk4,6 activities.

EXAMPLE 3: Identification of pluripotent and other stem cells

5 The successful isolation, maintenance in vitro, genetic manipulation and germ-line transmission of pluripotent cells from species other than mouse has generally been difficult and largely unsuccessful to date. One of the difficulties in obtaining pluripotent cells from other species has been the inability to demonstrate simply and rapidly, that putative pluripotent cells are pluripotent.

10 Generally Oct-4 expression has been identified as a reliable marker for pluripotency. Alkaline phosphatase has also been used, but its absolute reliability as a pluripotent cell marker is questionable since a number of differentiated cells also express AP. Recently some novel markers that distinguish between different types of pluripotent cells were identified.

15 (PCT/AU99/00265). The markers L17 and Psc1 are expressed in ES cells and are down regulated in EPL cells. Conversely K7 is upregulated in EPL cells compared to ES cells.

 We have found that the expression and activity of cell cycle markers can be determined rapidly in vitro, and this data can be used alone, or in

20 combination with other markers not directly involved with cell cycle regulation, to establish the pluripotent state or otherwise of putative pluripotent cell populations. In particular the type of cell cycle regulatory activities present,

- 51 -

cell cycle regulatory molecules expressed, and the phosphorylation state of pRb can be used as markers for pluripotency.

The following criteria can be used to identify pluripotent cells.

- Elevated levels and constitutive activity of cyclin E-Cdk2 activities, and
5 cyclin E expression.

- Elevated levels and constitutive activity of cyclin A-Cdk2 activities and cyclin A expression.

- Inactive/constitutively phosphorylated pRb and/or other pRb tumour suppressor family members, such as p107 and p130.

10 Substantial reduction in, or absence of, the INK Cdk inhibitor p16.

Substantial reduction in, or absence of, the Cdk inhibitors p21 and p27.

Many differentiated cells in the mammal are not replaced by proliferation of existing differentiated cells, but by the proliferation of precursor cells (multipotent stem cells). Cellular decisions taken by stem cells are
15 exquisitely regulated by external cues, which link stem cell behavior to the requirements of the organism. Rates of renewal and differentiation are coupled to prevent depletion of the stem cell population and control the rate of differentiated cell production, while the pathway of stem cell differentiation can be varied in response to environmental stimuli.

20 These cells combine a number of properties that are shared with pluripotent cells, and distinguish them from terminally differentiated somatic

20 A major problem in obtaining pluripotent cells from species other than

mouse has been the inability to proliferate and maintain pluripotent cells in vitro. Manipulation of cyclin activity or activities that can enforce high cyclin-Cdk activities in putative pluripotent cells, offers one approach to overcome these difficulties. These approaches would mimic/reproduce essential characteristics of the stem cell-state.

For example constitutive expression of cyclin E from a transgene expression construct would force continuing rounds of cell proliferation by maintaining pRb in an inactive/phosphorylated state, and would promote constitutive transcription of E2F target genes. It is also likely that such cell cycle regulatory molecules are associated with chromatin remodelling and maintenance of pluripotent status (Brehm & Kouzarides, 1999; Kouzarides 1999). Alternatively, maintaining high Cdk activity through the established properties of upstream regulators such as c-myc (Amati et al 1998) is another approach to stabilize pluripotent cells in vitro. Another approach would be to enforce the elevated activity of down-stream effector molecules associated with these pathways.

Differentiation of pluripotent cells is associated with a change in cell cycle structure, changes in the rate of cell proliferation, establishment of cell cycle regulated Cdk activities and down regulation in the absolute activities of some Cdks. Maintenance of Cdk activities in a constitutive state would serve to maintain stem cells, prevent them from differentiating and to facilitate their propagation in culture for extended periods.

Materials and Methods

c-myc expression construct:

A construct suitable for the stable expression of human c-myc was constructed by inserting a Eco RI fragment, from the plasmid pSR• MSVtkNEO (Sawyers et al 1992) that contains the entire c-myc coding sequence, into the Eco RI site of the expression vector, pEF-IRES-puro (Hobbs et al 1998). This construct, pc-Myc.puro, utilizes the EF1• promoter and drives the transcription of the c-myc open reading frame in addition to the puroR gene by way of an internal ribosome entry site (IRES).

Establishment of stable c-myc expressing cells lines:

10• g of Ase I- digested pc-Myc.puro was electroporated into 5 X 10⁶ D3 ES cells under the following conditions. Sub-confluent ES cells were trypsinized, washed once in PBS and resuspended in 900• l PBS (5.6 X 10⁷ cells/ml) with DNA. Cells/DNA were placed in a Bio Rad electroporation cuvette and electroporated at 500• F, 0.2KV. Cells were immediately resuspended in 10 ml complete DMEM in the absence of puromycin (Sigma). 0.6-3 x 10⁶ cells were then seeded into gelatinized 10cm diameter tissue culture grade petri dishes in 10 ml fresh DMEM plus LIF in the absence of puromycin. After 18 hours, media was replaced with fresh complete DMEM plus LIF and 1-5 • g/ml puromycin. Cells were selected for over a 10-14 day period. Media was changed during selection every 24 hours. Clonal cell lines were initially amplified in gelatinized 24 well plates (Falcon) amplified and

- 55 -

expression of human c-myc confirmed by Western blot analysis using the anti c-myc monoclonal antibody, 9E10 (Evan et al 1985) on crude whole cell lysates.

Preparation of whole cell protein extracts:

5 Whole cell extracts were prepared by washing cell pellets in ice cold PBS and resuspending in ice cold lysis buffer (250mM KCl, 50mM Hepes pH 7.9, 0.1mM EDTA, 0.1mM EGTA, 0.4mM NaF, 0.4mM NaVO₄, 10% glycerol, 0.1% Tween 20, 0.5mM PMSF, 1mg/ml leupepin, 1mM DTT) at 1.5×10^8 cells/ml. Protein concentrations of extracts were typically 5mg/ml.

LIF titration assay:

10 LIF titration assays were set up in gelatinized 24 well trays. 900 μ l of DMEM LIF was added to each well together with varying amounts of recombinant LIF were added to give final concentrations of 40-0 U/ml (ESGro, AMRAD). 500 cells were added to each well and the medium mixed for even
15 spreading. After 6 days the plates were stained for alkaline phosphatase activity.

Alkaline phosphatase staining:

Alkaline phosphatase was visualized using the diagnostic kit 86-R (Sigma). The kit was used according to the manufacturer's specifications with
20 the following modification; cell layers were fixed in 4.5 mM citric acid, 2.25 mM sodium citrate, 3 mM sodium chloride, 65% methanol and 4% para-

- 56 -

formaldehyde prior to washing and staining.

Results

c-myc drastically reduces spontaneous differentiation of ES cells

Examination of c-myc expressing colonies in comparison to control
5 (vector alone, puromycin-selected colonies) revealed striking differences. First,
stably-transfected c-myc ES cell lines were far more uniform in shape and size
(ie uniform dome shaped colonies; data not shown). Besides forming
spherical, domed-shaped colonies, a noticeable reduction in the number of
differentiated cells surrounding the colonies was observed. This is in marked
10 contrast to control ES cells which are prone to spontaneous differentiation (5-
10 % colonies on a plate normally exhibit a differentiated phenotype) in
addition to the presence of differentiated cells at the periphery of colonies.
Less than 1% of c-myc transfected cells exhibited a differentiated phenotype
(data not shown). These characteristics were consistent in all ES colonies that
15 expressed the c-myc transgene (12 c-myc and 12 control cell lines were
characterized).

c-myc reduces the requirement that ES cells have for LIF

To evaluate the stability of c-myc ES colonies and their ability to
differentiate, we compared the ability of these cells to retain pluripotency in the
20 presence of reduced levels of LIF. The logic being that if c-myc was promoting
pluripotency and blocking spontaneous differentiation, it may substitute,
partially or fully, for LIF's stem cell maintenance function. This possibility was

20 The ability of c-myc to maintain ES cell pluripotency was characterized in further detail using conditions of cell growth that would normally facilitate the differentiation of cells, resulting in loss of pluripotency. ES cells were converted into pluripotent EPL cells (see Example 1) and grown on bacterial

petri dishes as aggregate-forming EPL embryoid bodies in the absence of LIF (described in Rathjen et al 1999). The differentiation of control (vector minus c-myc insert) versus c-myc transfected cells was assessed by Northern blot analysis over an 5-day time course, using Oct4 (marker of pluripotency) and brachyury (marker for nascent mesoderm) probes. Figure 14 shows that vector control cell lines down-regulate Oct4 mRNA and lose pluripotency after 3 days (after LIF withdrawal) whereas all c-myc cell lines tested maintain high levels of Oct4 expression throughout the experiment, confirming that they retain pluripotency. In control cell lines, brachyury mRNA was markedly up-regulated at day 3, indicating the formation of nascent mesoderm. In c-myc cell lines, the expression of brachyury was typically delayed by at least a day and was not up-regulated to the same extent as in vector alone cell lines. This indicates that differentiation of c-myc expressing cells was incomplete and largely blocked. Differentiation that did occur was delayed significantly. Hence, in the absence of LIF, the differentiation of pluripotent cells is severely compromised by the activity of c-myc. This shows that c-myc has properties that allow for the stabilization maintenance of pluripotent stem cells.

Other parameters were also evaluated with regard to the inability of ES cells ectopically expressing c-myc to differentiate (and hence retain pluripotency), such as retention of high cyclin E expression levels and Cdk2-cyclin E activity. In c-myc expressing cell lines, cyclin E protein remained elevated to similar levels as seen in LIF-maintained ES cells (data not shown). Cdk2-cyclin E associated kinase activity decreased approximately 2-3 fold

Controlled differentiation is central to many of the commercial applications of pluripotent cells. For differentiation of the pluripotent cells with ectopic cell cycle regulatory activities, release from these activities is a major advantage. For this reason the use of protein transduction approach to introduce cell cycle regulatory molecules into the cell has significant advantages, since it allows activity to be delivered and persist for just the time required. Levels of cell cycle regulatory molecules, sufficient for effective activity, can be achieved for the time required. Once the activities have served

- 60 -

their purpose, intracellular levels of the transduced protein are reduced as cells replicate, and further reduction is achieved by normal protein degradation. Where cell cycle regulatory activity is achieved by ectopic gene expression, release could be achieved by use of a c-myc, cyclin E or other appropriate
5 expression constructs that includes Cre-lox recognition sequences. The expression construct would be removed from pluripotent cells by application of the Cre-lox system. Alternatively an inducible promoter could be included in the expression construct. Constitutive c-myc or cyclin E expression is achieved so long as the inducer is included in the culture medium, and when
10 differentiation of the pluripotent cells is required, the inducer would be removed.

Example 5: selection against differentiated prb+ cells using ectopic p16 expression: an approach to select for pluripotent stem cells

A system to allow and promote the maintenance of pluripotent stem
15 cells and select against differentiated cells is attractive for several reasons;

it will eliminate differentiated cells that can destabilise pluripotent stem cells, and thus improve conditions under which pluripotent cells can be maintained

(i) it can be used to functionally select for other types of stem cells
20 with similar properties to ES cells. Such properties would also be expected of other stem cells populations that have been difficult to identify, isolate and maintain in vitro.

- 61 -

Our characterization of pluripotent ES cells has shown that pRb is unable to associate with at least some of its cellular targets (specifically E2F transcription factors, see Example 2). The basis for this is likely to be due to the precocious activity of Cdks such as Cdk2-cyclin E and or Cdk2-cyclin A.

5 These cells can therefore be considered as having no functional pRb, as it is biochemically inactive. As stated previously (see Example 2), this eliminates the need for Cdk4-cyclin D activities and should hence render these cells insensitive to the Cdk inhibitory molecule, p16 (see Jiang et al 1988 and references therein). This marks what is a fundamental difference between the
10 cell cycle regulation of differentiated cells compared to pluripotent cells. This fundamental difference in cell cycle control mechanisms can be exploited for the selection purpose described above as pluripotent cell that do not utilize Cdk4,6-cyclin D activities will be insensitive to p16, whereas differentiated pRb+ cells will be sensitive (Jiang et al 1998).

15 **Materials and methods:**

All methods and materials were as described in previous examples except where stated

p16 expression construct and characterization of p16 cell lines:

A Xho I-Not I fragment from the plasmid pKS.mp16 (gift from C. Sherr)
20 spanning the entire coding region of human p16 was subcloned into the vector pEF-IRES (see Example 4). The resulting construct was linearized with Nde I and transfected into D3 ES cells as described previously (see Example 4). Puromycin resistant colonies (1-5 • g/ml puromycin) were cloned, amplified

and analyzed for p16 expression using an antibody raised against p16 (Santa Cruz sc-1207) by Western blot analysis (data not shown). The functionality of the p16 fusion protein was tested by its ability to bind Cdk4 in whole cell lysates. Typical cell lines quantitatively sequestered all Cdk4/Cdk6, indicating
 5 that all Cdk4/6 was inactive (data not shown). Differentiated cells under these conditions would exhibit a G1 arrest (Sherr and Roberts 1999). p16 is not normally expressed at detectable levels in ES cells (data not shown).

Selection against differentiated cells using transduced p16 protein

As another approach to determine that p16 can be used to select
 10 against differentiated cells, recombinant TAT-p16 fusion proteins were produced (as described in Nagahara 1998) in bacteria. Typically, recombinant TAT-p16 was purified from 200ml culture of cells grown to an OD₆₀₀ 1.0. Cells were collected by centrifugation and resuspended in 20 ml Buffer Z (8M urea, 100mM NaCl, 20mM Hepes pH 8.0). The cell lysate was sonicated for 15
 15 seconds, three times and centrifuged at 12,000 rpm for 15 minutes at 4°C. The supernatant was adjusted to 20mM imidazole and tumbled for 15 minutes in a 50 ml Falcon tube with 5 ml nickel-affinity resin (Qiagen) for purification of the 6X Histidine-tagged TAT-p16 fusion protein. The lysate and beads were poured into a 10ml Bio Rad EconoColumn and after further equilibration and
 20 elution of unbound protein with 50 ml Buffer Z, 20mM imidazole, TAT-p16 protein was eluted with 10 ml Buffer Z, 100mM imidazole (1ml fractions). Fractions containing the TAT-p16 fusion were identified by loading 5•l samples from each fraction on a 12% SDS polyacrylamide running gel,

Technology that allows reversion of differentiated cells to a pluripotent/stem cell state or for the selection of such events, has many

applications that include cell therapy approaches for the treatment of diseases, and the genetic manipulation of animals.

Several approaches have been developed to achieve reprogramming, including nuclear transfer technology. In recent developments adult cells have
5 been used as donors in nuclear transfer (Wilmut et al,1997), and has resulted in the production of cloned animals, albeit with very low efficiency. Reprogramming of bone marrow cells has also occurred following bone marrow transplantation, and cells derived from bone marrow have been identified in several non-haematopoietic tissues (Eglitz & Mezey, 1997; Shi et
10 al, 1998).

Despite improvements in the ability to reprogramme partially differentiated or differentiated cells, there are inadequacies with the current technology that restrict its widespread application for commercial, medical and agricultural benefit. Nuclear transfer, particularly with differentiated cells, is
15 very inefficient, and with all currently available technology it is not possible to identify or maintain cells in transitional states of dedifferentiation. Neither is it possible to direct the redifferentiation of reprogrammed cells in a controlled manner.

Major improvements in reprogramming technology are provided by
20 approaches that involve manipulation of cell cycle regulatory molecules.

One approach involves:

the elevation of cyclin E-Cdk2 activity and/or cyclin A-Cdk2 activity, either directly or by promoting the activities of upstream regulators of cyclin E-Cdk2 activity and cyclin A-Cdk2 activity such as c-myc, in differentiated cells. Elevation of these activities can be achieved by transformation of gene
5 expression constructs, or by transduction of cyclin E, Cdk2 and/or Cyclin A/Cdk2 and/or c-myc polypeptides, by methods known to promote polypeptide entry into cells. These methods include electroporation, lipofection, microinjection and use of fusion proteins comprised of a transduction domain such as included in the HIV peptide TAT (Nagahara et al, 1998).

10 Cells manipulated in this way may revert to a less differentiated state without further manipulation. Alternatively manipulation in this way may be used to "prime" cells so that they are more responsive to other reprogramming signals, such as the signals that operate during nuclear transfer, where the genetic information from differentiated or partially differentiated cells is
15 reprogrammed by transfer into an enucleated oocyte. Use of "primed" cells as nuclear transfer donors, where cell cycle activities more closely resemble those of dedifferentiated cells, would improve the efficiency of reprogramming. The nuclear transfer embryo formed from such "primed" cells may be allowed to develop, leading to the production of a live animal. Alternatively the
20 "primed" cells could be fused with a cytoplasm prepared from a pluripotent cell, thus leading to the production of pluripotent cells with the genetic characteristics of the donor-differentiated cell. Nuclear transfer products using both oocyte cytoplasm or pluripotent cell cytoplasm can be used as a source of

pluripotent cells for in vitro genetic manipulation and/or controlled differentiation.

A second approach involves:

the selection of dedifferentiated cells based on their insensitivity to the
5 Cdk inhibitor, p16 and other members of the INK family of Cdk inhibitors with
similar activity . For example, pluripotent cells can be 'captured' from a
differentiated cell population by selecting against differentiated cells with p16
fusion proteins, where p16 is linked to a transduction domain, such as the
transduction domain of the TAT protein. This approach, could be used in a
10 manner similar to that described in Example 5, in several scenarios including
the following. (1) where cells spontaneously dedifferentiate in vitro at low
frequency. (2) where cells are reprogrammed in response to inductive
environmental signals such as provided by enucleated oocytes, enucleated
pluripotent cells, inclusion with pluripotent cells in culture, or in differentiating
15 embryoid bodies.

Due to limitations with the current art, dedifferentiated cells formed
either spontaneously or in response to an inductive environment, cannot be
recognised or maintained in a dedifferentiated state. Use of the Cdk inhibitor
p16, or other members of the INK family of Cdk inhibitors with similar activity,
20 allows such dedifferentiated cells to be captured in vitro. Furthermore the
maintenance in an undifferentiated state of cells captured by p16 expression,
can be enhanced by expression of other cell cycle regulatory molecules such
as cyclin E-Cdk2, cyclin A-Cdk2, and c-myc, that enforce pluripotent cell
maintenance and proliferation in vitro (see Example 4).

It will be understood that the invention disclosed and defined in the specification extends to all alternative combinations of two or more of the individual features mentioned or evident from the text or drawings. All of these different combinations constitute various alternative aspects of the invention.

- 5 It will be understood that the term "comprises" or its grammatical variants as used herein is equivalent to the term "includes" and is not to be taken as excluding the presence of other elements or features.

Evan GI, Lewis GK, Ramsay G, Bishop JM. Isolation of monoclonal antibodies specific for human c-myc proto-oncogene product. (1985)

- Hall M & Peters G (1996). Genetic alterations of cyclins, cyclin-dependent kinases, and Cdk inhibitors in human cancer. *Adv Cancer Res* 68, 67-108
- 5 Hobbs S, Jitrapakdee S, Wallace JC. (1998). Development of a bicistronic vector driven by the human polypeptide chain elongation factor 1alpha promoter for creation of stable mammalian cell lines that express very high levels of recombinant proteins. *Biochem Biophys Res Commun* 252, 368-372.
- 10 Hogan B, Beddington R, Constantinti F & Lacy E (1994). *Manipulating the mouse embryo. A laboratory manual. Second edition.* Cold Spring Harbor Laboratory Press.
- 15 Hooper M, Hardy K, Handyside A, Hunter S & Monk M. (1987). HPRT deficient (Lesch-Nyhan) mouse embryos derived from germ-line colonization by cultured cells. *Nature* 326, 292-295.
- Janne J, Alhonen L, Hyttinen JM, Peura T, Tolvanen M & Korhonen VP (1998). Transgenic bioreactors. *Biotech Ann Rev* 4, 55-74
- 20 Jiang H, Chou HS & Zhu L (1998). Requirement of cyclin E-Cdk2 inhibition in p16^{INK4a}-mediated growth suppression. *Mol Cell Biol* 18, 5284-5290
- Kam J, Watson JV, Lowe AD, Green SM and Vedeckis W (1989). Regulation of cell cycle length by c-myc levels. *Oncogene*. 4, 773-787.
- 25 Kawasaki H, Eckner R, Yao TP, Taira K, Chiu R, Livingstone DM & Yokoyama KK (1998). Distinct roles of the co-activators p300 and CBP in retinoic-acid-induced F9-cell differentiation. *Nature* 393, 284-289

- Knowles BB, Pan S, Solter D, Linnerbach A, Croce C & Huebner K (1980). Expression of H-2, laminin, and SV40 T and TASA on differentiation of transformed murine teratocarcinoma cells. *Nature*. 288, 615-618.
- 5 Koh J, Enders GH, Dynlacht BD, Harlow E. (1995). Tumour-derived p16 alleles encoding proteins defective in cell-cycle inhibition. *Nature*. 375, 506-10.
- Kouzarides T (1999). Histone acetylases and deacetylases in cell proliferation. *Curr Opin Genet Dev* 9, 40-48
- 10 Nagahara H, Vocero-Akbani AM, Snyder EL, Ho A, Latham DG, Lissy NA, Becker-Hapak M, Ezhevsky SA, Dowdy SF (1998). Transduction of full-length TAT fusion proteins into mammalian cells: TAT-p27Kip1 induces cell migration. *Nat Med*. 4, 1449-52.
- Pagano M, Pepperkok R, Verde F, Ansorge W & Draetta G (1992). Cyclin A is required at two points in the human cell cycle. *EMBO J* 11, 961-971
- 15 Palermo T & Peters G (1996). Perturbation of cell cycle regulators in human cancer. *Cancer Surv* 27, 351-367
- Pease S, Braghetta P, Gearing D, Grail D & Williams RL (1990). Isolation of embryonic stem (ES) cells in media supplemented with recombinant leukemia inhibitory factor (LIF). *Dev. Biol.* 141, 344-352.
- 20 Platt JL & Lin SS (1998). The future promises of xenotransplantation. *Ann NY Acad Sci* 862, 5-18
- Rathjen PD, Lake J, Whyatt LM, Bettles MD & Rathjen J (1998). Properties and uses of embryonic stem cells: prospects for application
- 25

to human biology and gene therapy. Reprod Fert Dev 10, 1-17

Rathjen J Lake JA, Bettess MD, Washington JM, Chapman G & Rathjen PD (1999). Formation of a primitive ectoderm like cell population, EPL cells, from ES cells in response to biologically derived factor. J Cell Sci. 112, 601-612.

Resnitzky D, Gosen M, Bujard H & Reed SI (1994). Acceleration of the G1/S transition by expression of cyclins D1 and E with an inducible system. Mol. Cell. Biol. 14, 1669-1679.

Sambrook et al (1989) Molecular cloning, A Laboratory Manual, Second Edition, Cold Spring Harbour Laboratory Press, Cold Spring Harbour, New York

Sawyers CL, Callahan W, Witte ON (1992). Dominant negative MYC blocks transformation by ABL oncogenes. Cell. 70, 901-10.

Sherr CJ (1994). G1 phase progression: cycling on cue. Cell. 79, 551-555.

Sherr CJ & Roberts JM (1995).). Inhibitors of mammalian G1 cyclin-dependent kinases. Genes and Devel 9, 1149-1163

Sherr CJ, and Roberts JM (1999). CDK inhibitors: positive and negative regulators of G1-phase progression. Genes Dev.13, 1501-12

Shi Q, Raffii S, Wu MH, Wijelath ES, Yu C, Ishida A, Fujita Y, Kothari S, Mohle R, Sauvage R et al, (1998). Blood 92, 362-367

Smith AG (1991). Culture and differentiation of embryonic stem cells. J. Tiss. Cult. Meth. 13, 89-94.

5

1. A method of regulating the mitotic and/or physiological activities, and differentiation potential of a pluripotent or multipotent cell, which method includes manipulating the expression and/or activity of a cell cycle regulatory molecule including a regulatory molecule selected from one or more of the groups consisting of a cyclin, a cyclin-dependent protein kinase (Cdk), a Cdk inhibitor, upstream regulators thereof or biochemical targets thereof and/or tumour suppressor protein, and molecules displaying similar activities, in a pluripotent or multipotent cell.

2. A method according to claim 1 wherein the activity of the cell cycle regulatory molecule is manipulated by increasing or decreasing the level of expression of said molecules.

3. A method according to claim 1 wherein the pluripotent cells or multipotent cells are selected from one or more of a group consisting of epiblast cells, embryonic stem (ES) cells, primitive ectoderm-like (EPL) cells, primordial germ cells (PGCs) or embryonic carcinoma (EC) cells.

4. A method according to claim 1 wherein the cell cycle regulatory molecule includes a cyclin selected from one or more of the group consisting of cyclin D, cyclin E and cyclin A or a molecule exhibiting similar activity or a functionally active fragment or analogue thereof.

5. A method according to claim 1 wherein the cell cycle regulatory molecule includes a cyclin-dependent protein kinase (Cdk) selected from one or more of the group consisting of Cdk4, Cdk6 or Cdk2 or a molecule exhibiting similar activity or a functionally active fragment or analogue thereof.

6. A method according to claim 1 wherein the cell cycle regulatory molecule includes a Cdk inhibitor selected from the INK, CIP or KIP families.

- 75 -

7. A method according to claim 6 wherein the Cdk inhibitor is selected from one or more of the group consisting of p27, p57, p16, p15, p18, p19 or p21 or a molecule exhibiting similar activity or a functionally active fragment or analogue thereof.

5 8. A method according to claim 1 wherein the cell cycle regulatory molecule includes a tumour suppressor protein selected from one or more of the group consisting of retinoblastoma protein (pRb), p107 or p130 or a molecule exhibiting similar activity or a functionally active fragment or analogue thereof.

10 9. A method according to claim 8 wherein the activity or phosphorylation state of the tumour suppressor protein is manipulated.

10. A method according to claim 1 wherein the upstream regulator is selected from the group consisting of one or more of proto-oncogenes *myc* and *ras* and upstream signalling pathways Raf, MAP Kinase or Rho.

15 11. A method according to claim 10 wherein the upstream regulator is *myc* or *ras* and the activity of the regulator is increased.

12. A method for identifying pluripotent or pluripotent-related cells which method includes analysing the cell population for pluripotent cell cycle characteristics including one or more of pluripotent-specific cell cycle
20 structure; pluripotent-specific expression of cell cycle regulatory molecules; and phosphorylation status of a tumour-suppressor protein(s).

13. A method according to claim 12, wherein the method includes measuring expression of cell cycle regulatory molecules, including a cyclin(s), a cyclin-dependent protein kinase (Cdk), a Cdk inhibitor, upstream regulators
25 of said molecules or biochemical targets thereof.

14. A method according to claim 13, wherein the pluripotent or

manipulating the expression and/or activity of a cell cycle regulatory molecule, including a regulatory molecule selected from one or more of the groups consisting of a cyclin, a cyclin-dependent protein kinase (Cdk), a Cdk inhibitor, upstream regulators thereof or biochemical targets thereof and/or

- 77 -

tumour suppressor protein, and molecules displaying similar activities, in a pluripotent or multipotent cell such that some or all of the features of the cell cycle of pluripotent cells are enforced.

20. A method according to claim 19 wherein constitutive cyclin E
5 activity and/or constitutive cyclin A activity is increased.

21. A method according to claim 19 wherein constitutive cyclin E activity and/or cyclin A activity is increased by enforcing expression of exogenous cyclin E and/or cyclin A activity, and/or enforcing expression of Cdk
2.

10 22. A method according to claim 19 wherein the level of constitutive cyclin E activity and/or cyclin A activity is increased utilising the proto-oncogenes *myc* and *ras*.

23. A method according to claim 19 wherein constitutive cyclin D activity is increased by expression of exogenous cyclin D or expression of Cdk
15 4 and/or Cdk 6.

24. A method according to claim 19 wherein tumour suppressor proteins are inactivated such that maintenance and proliferation of pluripotent cells is prolonged.

25. A method according to claim 24 wherein the tumour suppressor
20 protein pRb is inactivated utilising hyperphosphorylation, antisense techniques or gene inactivation.

26. A method according to claim 19 wherein the upstream regulator is selected from the group consisting of

one or more of proto-oncogenes *myc* and *ras* and upstream
25 signalling pathways Raf, MAP Kinase or Rho.

34. A method according to claim 33 wherein the upstream regulator is myc or ras and the activity of the regulator is increased.

multipotent cells.

42. A method according to claim 35 wherein the multipotent or pluripotent cells are cells derived by reversion from a differentiated or partially differentiated state to a less differentiated state.

5 43. A method according to claim 42 wherein the reversion is spontaneous or environmentally induced.

44. A method according to claim 35 wherein the cells are in a transitional pluripotent state.

10 45. A method according to claim 42 which method includes initiating ectopic proto-oncogene expression and selecting for cells exhibiting survival and maintenance.

46. A method according to claim 45 wherein the proto-oncogene is *myc*.

15 47. A method according to claim 35 wherein the maintenance and/or proliferation of the selected pluripotent or multipotent cells is facilitated by further manipulating the activity of a cell cycle regulatory molecule.

20 48. A method of regulating the differentiation of pluripotent or multipotent cells which method includes manipulating the expression and/or activity of a cell cycle regulatory molecule, including a regulatory molecule selected from one or more of the groups consisting of a cyclin, a cyclin-dependent protein kinase (Cdk), a Cdk inhibitor, upstream regulators thereof or biochemical targets thereof and/or tumour suppressor protein, and molecules displaying similar activities, in a pluripotent or multipotent cell, in said cells.

25 49. A method according to claim 48, wherein the differentiation of the pluripotent or multipotent cells is regulated by one or more of the following

manipulations:

increasing the activity of the tumour suppressor protein;
reducing cyclin E and/or cyclin A-associated activities;
the activity of Cdk2 becomes cell cycle regulated; or
5 upregulating the activities and/or expression of Cdk inhibitors.

50. A method of regulating the cell cycle of primary or untransformed cells, which method includes

manipulating the expression and/or activity of a cell cycle regulatory molecule, such that the proliferation and maintenance of
10 differentiated cells are reduced, wherein the regulatory molecule is selected from one or more of the groups consisting of a cyclin, a cyclin-dependent protein kinase (Cdk), a Cdk inhibitor, upstream regulators thereof or biochemical targets thereof and/or tumour suppressor protein, and molecules displaying similar activities, in a primary or untransformed cell.

15 51. A method according to claim 50 wherein the lifespan of the differentiated or partially differentiated cells is prolonged by one or more of

upregulation of Cyclin E-, Cyclin D- or Cyclin A-associated activities, or upregulation of upstream regulators of cyclin-associated activities.

20 52. A method of identifying differentiating cells in a mixed cell population, which method includes analysing the cell population for differentiating cell cycle characteristics including

differentiation-specific cell cycle structure;
differentiation-specific expression and/or
activity of cell cycle regulatory molecules
25 the presence of active tumour suppressor proteins.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 April 2001 (05.04.2001)

PCT

(10) International Publication Number
WO 01/23531 A1

(51) International Patent Classification⁷: C12N 5/08, 5/06

(21) International Application Number: PCT/AU00/01184

(22) International Filing Date:
22 September 2000 (22.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PQ 3073 24 September 1999 (24.09.1999) AU

(71) Applicant (for all designated States except US): LUMI-
NIS PTY LTD [AU/AU]; 1st Floor, 10-20 Pulteney Street,
Adelaide, S.A. 5000 (AU).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RATHJEN, Peter,
David [AU/AU]; 1 Mimosa Avenue, Blackwood, S.A.
5051 (AU). DALTON, Stephen [AU/AU]; 43 Carlisle
Road, Westbourne Park, S.A. 5041 (AU).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— With international search report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: RECOGNITION OF DIFFERENCES IN CELL CYCLE STRUCTURE BETWEEN STEM AND DIFFERENTIATED
CELLS

(57) Abstract: The invention resides in the recognition of differences in cell cycle structure between stem and differentiated cells. Pluripotent cells spend ~65 % of time in S phase, ~15 % in G1 and ~20 % in M. Partially differentiated cells have a different cell cycle structure, e.g. mesoderm spend 15-20 % of time in S phase, and 60 % in G1. These differences in cell cycle have lead to the identification of differences in expression of cell cycle genes and activity of regulators. The claims are directed to manipulating cell cycles genes, regulators and proteins to regulate cell activity or cells cycle, identify cell types, facilitate maintenance, reprogramme cells, or regulate differentiation.

WO 01/23531 A1



1. Regulated activation of pRb-kinases (Cdk)

2. pRb phosphorylation

3. Derepression of genes required for G1-S progression

- 1. no regulated pRb kinase (Cdk2-cyclinE)
- 2. pRb constitutively hyper-phosphorylated
- 3. E2F target genes constitutively active

Normal somatic cells

Embryonic cells
(tumour cells)

Figure 1

Remodelling of the cell cycle during embryonic development/differentiation






Figure 2

Figure 3



Early mouse embryogenesis. A. Schematic representation of mouse embryogenesis, between 4.5 and 7.0 d.p.c., highlighting the pluripotent cell populations and prior to and during the onset of gastrulation. Gastrulation initiates at the primitive streak and results, initially, in the formation of mesoderm. B. Pluripotent cells from 6.5 d.p.c. embryos were isolated, labeled with propidium iodide and subjected to flow cytometry analysis.

Figure 4



Epiblast Cells (4.4 hours)

G1	S	G2/M
----	---	------

Embryonic Stem Cells (10 hours)

G1	S	G2/M
----	---	------

EPL Cells (8-10 hours)

G1	S	G2/M
----	---	------

Balb/c 3T3 Fibroblast Cells (24 hours)

G1	S	G2/M
----	---	------

Figure 5
Cell cycle remodelling during differentiation of cells in embryoid bodies






Figure 6

E2F target genes are not cell cycle regulated in ES cells




Cell cycle regulation of E2F transcripts in mouse fibroblasts



WO 01/23531

PCT/AU00/01184

Figure 7



E2F complexes in pluripotent cells are predominantly free of pRb family members

E2F4 is the major E2F
activity in ES cells




Figure 8




Figure 10





Figure 11



Cyclin E and cyclin A associated kinase activities collapse during differentiation




Figure 12








Figure 15





Attorney Docket No.:

Title: Recognition of differences in cell cycle structure between stem and differentiated cells

Page 1

(Foreign associate use only)

DECLARATION AND POWER OF ATTORNEY

Attorney's Docket No. _____

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: Recognition of differences in cell cycle structure between stem and differentiated cells, the specification of which is attached hereto.

was filed on 22 September 2000 as U.S. Application or PCT International Application No. PCT/AU00/01184 and was amended (if applicable) on _____.

to the
best of my
knowledge

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above. I do not know and do not believe that the same was ever known or used by others in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to the date of this application. I further state that the invention was not in public use or on sale in the United States of America more than one year prior to the date of this application. I understand that I have a duty of candor and good faith toward the Patent and Trademark Office, and I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign application for patent or inventor's certificate disclosing subject matter in common with the above-identified specification and having a filing date before that of the application on which priority is claimed:

Country	App. No.	Date of Filing	Priority Claimed Under 35 USC §119
Australia	PQ3073	24 September 1999	Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>
PCT	PCT/AU00/01184	22 September 2000	Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>

I hereby claim the benefit under Title 35, United States Code, § 120 of any prior United States application(s), or §365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each claim of the present application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations §1.53, which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Filing Date	Status: patented, pending, abandoned

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patents issuing thereon.



Attorney Docket No.:

Title: Recognition of differences in cell cycle structure between stem and differentiated cells

Page 1

(Foreign associate use only)

DECLARATION AND POWER OF ATTORNEY

Attorney's Docket No. _____

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: Recognition of differences in cell cycle structure between stem and differentiated cells, the specification of which

is attached hereto.

was filed on 22 September 2000 as U.S. Application or PCT International Application No. PCT/AU00/01184 and was amended (if applicable) on _____.

to the
best of my
knowledge

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above. I do not know and do not believe that the same was ever known or used by others in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to the date of this application. I further state that the invention was not in public use or on sale in the United States of America more than one year prior to the date of this application. I understand that I have a duty of candor and good faith toward the Patent and Trademark Office, and I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign application for patent or inventor's certificate disclosing subject matter in common with the above-identified specification and having a filing date before that of the application on which priority is claimed:

Country	App. No.	Date of Filing	Priority Claimed Under 35 USC §119
Australia	99/01184	24 September 1999	Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>
PCT	AU00/01184	22 September 2000	Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>

I hereby claim benefit under Title 35, United States Code, § 120 of any prior United States application or §365(c) of any PCT international application designating the United States of America listed below and, insofar as the subject matter of each claim of the present application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations §1.56, which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Filing Date	Status: patented, pending, abandoned
-----------------	-------------	--------------------------------------

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patents issuing thereon.

Attorney Docket No.:

Title: Recognition of differences in cell cycle structure between stem and differentiated cells

Page 2

I hereby authorize the U.S. attorneys named herein to accept and follow instructions from _____, as to any action to be taken in the Patent and Trademark Office regarding this application, without direct communication between the U.S. attorney and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney named herein will be notified by the undersigned.

POWER OF ATTORNEY: The following attorneys are hereby appointed to prosecute this application and transact all business in the Patent and Trademark Office connected therewith: Peter G. Pappas - 33,205; Daniel J. Warren - 34,272; William L. Warren - 36,714; Michael S. Pavento - 42,985; Lisa M. Coburn - 44,669; Robert A. Lester - 24,004; Erinn C. Kelly - 44,822; Jerry C. Liu - P47,754; Kevin W. King - 42,737.

Send correspondence to: SUTHERLAND ASBILL & BRENNAN LLP

Direct telephone calls at (404) 853-8000 to _____

999 Peachtree Street, NE

Atlanta, GA 30309-3996

Full name of sole or first inventor: Peter David RATHJEN	
Citizenship: Australian	
Residence: 1 Mimosa Avenue, South Australia 5051, Australia	
Post Office Address: "as above"	
Inventor's signature <i>P.D.R.</i>	Date: 02-08-02

Peter Rathjen

Full name of second inventor: Stephen DALTON	
Citizenship: Australian	
Residence: 43 Carlisle Road, Westbourne Park, South Australia, 5041, Australia	
Post Office Address: "as above"	
Inventor's signature <i>SD</i>	Date:

Steve Dalton

Attorney Docket No.:

Title: Recognition of differences in cell cycle structure between stem and differentiated cells

Page 2

I hereby authorize the U.S. attorneys named herein to accept and follow instructions from _____ as to any action to be taken in the Patent and Trademark Office regarding this application, without direct communication between the U.S. attorney and the undersigned. In the event of a change in the persons from whom instructions may be taken, the U.S. attorney named herein will be notified by the undersigned.

POWER OF ATTORNEY: The following attorneys are hereby appointed to prosecute this application and transact all business in the Patent and Trademark Office connected therewith: Peter G. Pappas - 33,205; Daniel J. Warren - 34,272; William L. Warren - 36,714; Michael S. Pavento - 42,985; Lisa M. Cobern - 44,669; Robert A. Lester - 24,004; Erinn C. Kelly - 44,822; Jerry C. Liu - P47,754; Kevin W. King - 42,737.

Send correspondence to: **SUTHERLAND ASBILL & BRENNAN LLP**

Direct telephone calls at (404) 853-8000 to _____


999 Peachtree Street, NE

Atlanta, GA 30309-3996

Full name of sole or first inventor: Peter David RATHJEN	
Citizenship: Australian	
Residence: 1 Mimosa Avenue, South Australia 5051, Australia	
Post Office Address: "as above"	
Inventor's signature <i>P.D.R.</i>	Date:

Full name of second inventor: Stephen DALTON	
Citizenship: Australian	
Residence: 43 Carlisle Road, Westbourne Park, South Australia, 5041, Australia	
Post Office Address: "as above"	
Inventor's signature <i>Stephen Dalton</i>	Date: 14/8/02

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not
present
for scanning. (Document title)

☒ Scanned copy is best available. *Some drawings are dark.*